

CONR¹⁰R¹¹, —CH₂R⁷, —CH₂R⁹, OR⁷, OR⁸ or OR⁹; or (C₁-C₆)-alkyl which is substituted by one or more radicals selected from the group consisting of halogen, (C₁-C₆)-alkoxy, (C₁-C₆)-haloalkoxy, (C₃-C₇)-cycloalkyl, S(O)_pR⁸, CO₂—(C₁-C₆)-alkyl, —O(C=O)—(C₁-C₆)-alkyl, NR¹⁰COR¹², NR¹⁰R¹¹, CONR¹⁰R¹¹, SO₂NR¹⁰R¹¹, OH, CN, N₂, OR⁷, NR¹⁰SO₂R⁸, COR⁸ and OR⁹;

A is (C₁-C₁₂)-alkylene and (C₁-C₁₂)-haloalkylene in which 2, 3 or 4 adjacent carbon atoms optionally form part of a (C₃-C₈)-cycloalkyl ring which is unsubstituted or substituted by one or more radicals selected from the group consisting of (C₁-C₆)-alkyl and halogen;

R⁵ is H, (C₃-C₆)-alkenyl, (C₃-C₆)-haloalkenyl, (C₃-C₆)-alkynyl, (C₃-C₆)-haloalkynyl, (C₃-C₇)-cycloalkyl, —(CH₂)_qR⁷, —(CH₂)_qR⁹ or NR¹⁰R¹¹ provided that for the last mentioned radical m is 2; or is (C₁-C₆)-alkyl unsubstituted or substituted by one or more radicals selected from the group consisting of halogen, (C₁-C₆)-alkoxy, (C₁-C₆)-haloalkoxy, (C₃-C₆)-alkenyloxy, (C₃-C₆)-haloalkenyloxy, (C₃-C₆)-alkynyloxy, (C₃-C₆)-haloalkynyloxy, (C₃-C₇)-cycloalkyl, S(O)_pR⁸, CN, NO₂, OH, COR¹⁰, NR¹⁰COR¹², NR¹⁰SO₂R⁸, CONR¹⁰R¹¹, NR¹⁰R¹¹, S(O)_pR⁷, S(O)_pR⁹, OR⁷, OR⁹ and CO₂R¹⁰;

R⁶ is (C₁-C₆)-alkyl, (C₁-C₆)-haloalkyl, (C₂-C₆)-alkenyl, (C₂-C₆)-haloalkenyl, (C₂-C₆)-alkynyl or (C₂-C₆)-haloalkynyl;

R⁷ is phenyl unsubstituted or substituted by one or more radicals selected from the group consisting of halogen, (C₁-C₆)-alkyl, (C₁-C₆)-haloalkyl, (C₁-C₆)-alkoxy, (C₁-C₆)-haloalkoxy, CN, NO₂, S(O)_pR⁸, COR¹¹, COR¹³, CONR¹⁰R¹¹, SO₂NR¹⁰OR¹¹, NR¹⁰OR¹¹, OH, SO₃H and (C₁-C₆)-alkylideneimino;

R⁸ is (C₁-C₆)-alkyl or (C₁-C₆)-haloalkyl;

R⁹ is heterocyclyl unsubstituted or substituted by one or more radicals selected from the group consisting of halogen, (C₁-C₄)-alkyl, (C₁-C₄)-haloalkyl, (C₁-C₄)-alkoxy, (C₁-C₄)-haloalkoxy, NO₂, CN, CO₂(C₁-C₆)-alkyl, S(O)_pR⁸, OH and oxo;

R¹⁰ and R¹² are each independently H, (C₁-C₆)-alkyl, (C₁-C₆)-haloalkyl, (C₃-C₆)-alkenyl, (C₃-C₆)-haloalkenyl, (C₃-C₆)-alkynyl, (C₃-C₆)-haloalkynyl, (C₃-C₆)-cycloalkyl, —(C₁-C₆)-alkyl-(C₃-C₆)-cycloalkyl, —(CH₂)_qR¹³ or —(CH₂)_qR⁹; or

R¹⁰ and R¹¹ and/or R¹⁰ and R¹² each together with the respective attached N atom form a five- or six-membered saturated ring which optionally contains an additional hetero atom in the ring

which is selected from O, S and N the ring being unsubstituted or substituted by one or more radicals selected from the group consisting of halogen, (C₁-C₆)-alkyl and (C₁-C₆)-haloalkyl; R¹¹ and R¹⁴ are each independently H, (C₁-C₆)-alkyl, (C₁-C₆)-haloalkyl, (C₃-C₆)-cycloalkyl or —(C₁-C₆)-alkyl-(C₃-C₆)-cycloalkyl;

R¹³ is phenyl unsubstituted or substituted by one or more radicals selected from the group consisting of halogen, (C₁-C₆)-alkyl, (C₁-C₆)-haloalkyl, (C₁-C₆)-alkoxy, (C₁-C₆)-haloalkoxy, CN, NO₂, S(O)_pR⁸ and NR¹¹R¹⁴;

R¹⁵ is R¹¹ or —(CH₂)_qR¹³;

m, n and p are each independently zero, one or two;

q is zero or one; and

each heterocyclyl in the above-mentioned radicals is independently a heterocyclic radical having 3 to 7 ring atoms and 1, 2 or 3 hetero atoms in the ring selected from the group consisting of N, O and S; or a pesticidally acceptable salt thereof.

5. (Previously Presented) 5-Substituted-alkylaminopyrazole derivatives of formula (I) as in claim 4, or pesticidally acceptable salts thereof, wherein:

R¹ is CN;

W is C-halogen or C—CH₃;

R² is hydrogen, halogen or CH₃;

R³ is (C₁-C₃)-haloalkyl, (C₁-C₃)-haloalkoxy or S(O)_p—(C₁-C₃)-haloalkyl;

R⁴ is hydrogen, (C₁-C₆)-alkyl or COR⁸;

A is (C₁-C₁₂)-alkylene and (C₁-C₁₂)-haloalkylene in which 2, 3 or 4 adjacent carbon atoms optionally form part of a (C₃-C₈)-cycloalkyl ring which is unsubstituted or substituted by one or more radicals selected from the group consisting of (C₁-C₆)-alkyl and halogen;

R⁵ is H, (C₃-C₆)-alkenyl, (C₃-C₆)-haloalkenyl, (C₃-C₆)-alkynyl, (C₃-C₆)-haloalkynyl, (C₃-C₇)-cycloalkyl, —(CH₂)_qR⁷, —(CH₂)_qR⁹ or NR¹⁰R¹¹ provided that for the last mentioned radical S(O)_m is SO₂; or is (C₁-C₆)-alkyl substituted by one or more radicals selected from the group consisting of halogen, (C₁-C₆)-alkoxy, (C₁-C₆)-haloalkoxy, (C₃-C₆)-alkenyloxy, (C₃-C₆)-haloalkenyloxy, (C₃-C₆)-alkynyloxy, (C₃-C₆)-haloalkynyloxy, (C₃-C₇)-cycloalkyl, S(O)_pR⁸,

CN, NO₂, OH, COR¹⁰, NR¹⁰COR¹², NR¹⁰SO₂R⁸, CONR¹⁰R¹¹, NR¹⁰R¹¹, S(O)_pR⁷, S(O)_pR⁹, OR⁷, OR⁹ and CO₂R¹⁰;

R⁶ is (C₁-C₆)-alkyl, (C₁-C₆)-haloalkyl, (C₂-C₆)-alkenyl, (C₂-C₆)-haloalkenyl, (C₂-C₆)-alkynyl or (C₂-C₆)-haloalkynyl;

R⁷ is phenyl unsubstituted or substituted by one or more radicals selected from the group consisting of halogen, (C₁-C₆)-alkyl, (C₁-C₆)-haloalkyl, (C₁-C₆)-alkoxy, (C₁-C₆)-haloalkoxy, CN, NO₂, S(O)_pR⁸, COR¹¹, COR¹³, CONR¹⁰R¹¹, SO₂NR¹⁰R¹¹, NR¹⁰R¹¹, OH, SO₃H and (C₁-C₆)-alkylideneimino;

R⁸ is (C₁-C₆)-alkyl or (C₁-C₆)-haloalkyl;

R⁹ is heterocyclyl unsubstituted or substituted by one or more radicals selected from the group consisting of halogen, (C₁-C₄)-alkyl, (C₁-C₄)-haloalkyl, (C₁-C₄)-alkoxy, (C₁-C₄)-haloalkoxy, NO₂, CN, CO₂(C₁-C₆)-alkyl, S(O)_pR⁸, OH and oxo;

R¹⁰ and R¹² are each independently H, (C₁-C₆)-alkyl, (C₁-C₆)-haloalkyl, (C₃-C₆)-alkenyl, (C₃-C₆)-haloalkenyl, (C₃-C₆)-alkynyl, (C₃-C₆)-haloalkynyl, (C₃-C₆)-cycloalkyl, —(C₁-C₆)-alkyl-(C₃-C₆)-cycloalkyl, —(CH₂)_qR¹³ or CH₂)_qR⁹; or

R¹⁰ and R¹¹ and/or R¹⁰ and R¹² each together with the respective attached N atom form a five- or six-membered saturated ring which optionally contains an additional hetero atom in the ring which is selected from O, S and N, the ring being unsubstituted or substituted by one or more radicals selected from the group consisting of halogen, (C₁-C₆)-alkyl and (C₁-C₆)-haloalkyl;

R¹¹ and R¹⁴ are each independently H, (C₁-C₆)-alkyl, (C₁-C₆)-haloalkyl, (C₃-C₆)-cycloalkyl or —(C₁-C₆)-alkyl-(C₃-C₆)-cycloalkyl;

R¹³ is phenyl unsubstituted or substituted by one or more radicals selected from the group consisting of halogen, (C₁-C₆)-alkyl, (C₁-C₆)-haloalkyl, (C₁-C₆)-alkoxy, (C₁-C₆)-haloalkoxy, CN, NO₂, S(O)_pR⁸ and NR¹¹R¹⁴;

R¹⁵ is R¹¹ or —(CH₂)_qR¹³;

m, n and p are each independently zero, one or two;

q is zero or one; and

each heterocyclyl in the above-mentioned radicals is independently a heterocyclic radical having 3 to 7 ring atoms and 1, 2 or 3 hetero atoms in the ring selected from the group consisting of N, O and S.

6. (Previously Presented) 5-Substituted-alkylaminopyrazole derivatives of formula (I) as in claim 4, or pesticidally acceptable salts thereof, wherein the symbols and indices in formula (I) have the following meanings:

R¹ is CN;

R² is chlorine;

R³ is CF₃ or OCF₃;

W is C—Cl;

R⁴ is hydrogen or (C₁-C₆)-alkyl;

R⁵ is (C₁-C₆)-alkyl;

R⁶ is CF₃;

A is (C₂-C₃)-alkylene

and m and n are each independently zero, one or two.

7. (Canceled).

8. (Canceled).

9. (Previously Presented) A pesticidal composition comprising a compound of formula (I) or a pesticidally acceptable salt thereof as defined in any one of claims 4 to 6, in association with a pesticidally acceptable diluent or carrier and/or surface active agent.

10. (Canceled)